

Discovery of dumbbell-shaped Cs^*He_n exciplexes in solid ^4He

D. Nettels, A. Hofer, P. Moroshkin,* R. Müller-Siebert, S. Ulzega, and A. Weis

*Département de Physique, Université de Fribourg,
Chemin du Musée 3, 1700 Fribourg, Switzerland†*

(Dated: February 9, 2008)

We have observed several new spectral features in the fluorescence of cesium atoms implanted in the hcp phase of solid helium following laser excitation to the 6^2P states. Based on calculations of the emission spectra using semiempirical Cs-He pair potentials the newly discovered lines can be assigned to the decay of specific Cs^*He_n exciplexes: an apple-shaped $\text{Cs}(\text{A}\Pi_{3/2})\text{He}_2$ and a dumbbell-shaped $\text{Cs}(\text{A}\Pi_{1/2})\text{He}_n$ exciplex with a well defined number n of bound helium atoms. While the former has been observed in other environments, it was commonly believed that exciplexes with $n > 2$ might not exist. The calculations suggest $\text{Cs}(\text{A}\Pi_{1/2})\text{He}_6$ to be the most probable candidate for that exciplex, in which the helium atoms are arranged on a ring around the waist of the dumbbell shaped electronic density distribution of the cesium atom.

PACS numbers: 76.70.Hb, 32.80.Wr, 32.30.Dx, 32.60.+i

Alkali atoms and helium atoms in their ground states strongly repel each other by virtue of the Pauli principle. However, an alkali atom excited to one of its P states can exert an attractive potential on a helium atom that can lead to bound states, known as exciplexes. The formation of alkali-helium exciplexes was considered for the first time by Dupont-Roc [1] and Karnorsky et al. [2] as an explanation for the observed quenching of atomic fluorescence from light alkali atoms (Na, Li) embedded in liquid or solid ^4He . In the meantime such molecules have been observed in different environments, such as liquid helium and cold helium gas [3, 4], as well as on the surface of helium nanodroplets [5, 6, 7, 8]. Here we present the first observations of such exciplexes in a solid helium matrix.

In earlier experiments [2] we have studied the excitation and fluorescence spectra of atomic cesium implanted into the bcc and hcp phases of solid helium. It was found that the excitation at the D_1 transition ($6\text{S}_{1/2}$ - $6\text{P}_{1/2}$) results in atomic fluorescence at the same transition, blue shifted (with respect to the free Cs atom) by the interaction with the helium matrix. At the same time, excitation on the D_2 transition ($6\text{S}_{1/2}$ - $6\text{P}_{3/2}$) produced merely a weak fluorescence on the D_1 emission line, which indicates that the $6\text{P}_{3/2}$ atoms are partly quenched into the $6\text{P}_{1/2}$ state, while the main relaxation channel remained unknown. Recently, the extension of the spectral range of our detection system allowed us to discover several new emission lines, red shifted with respect to the atomic fluorescence line. We attribute those lines to the formation and decay of Cs^*He_n molecules.

The maximal number of helium atoms for different alkali-helium exciplexes was found previously to be $n_{\text{max}} = 4$ for K^*He_n [8], $n_{\text{max}} = 6$ for Rb^*He_n [4], $n_{\text{max}} = 2$ for Cs^*He_n [3]. Hirano et al. [3] dis-

cuss on the basis of Cs^*He_2 -He potential energy surfaces that there should be no stable Cs^*He_3 configuration. They therefore conclude that exciplexes with more than $n_{\text{max}} = 2$ do not exist, since they regard the exciplex formation as a sequential process ($\text{Cs}^* \rightarrow \text{Cs}^*\text{He}_1 \rightarrow \dots \rightarrow \text{Cs}^*\text{He}_{n_{\text{max}}}$). However, our experimental results demonstrate unambiguously that in the hcp phase of solid helium $\text{Cs}(\text{A}\Pi_{1/2})\text{He}_n$ exciplexes with $n > 2$ are formed, when the cesium atoms are excited to the $6\text{P}_{3/2}$ state. From the relative integrated observed line intensities we conclude that the formation of those exciplexes is the most probable deexcitation channel of the $6\text{P}_{3/2}$ state.

In the present experiment a ^4He matrix doped with Cs atoms was produced by the technique described in our earlier papers [9, 10]. Data were taken in the hcp phase of solid ^4He at a temperature of 1.5 K and a pressure of 31.6 bar. For the excitation of the embedded atoms we used a single mode cw $\text{Ti}:\text{Al}_2\text{O}_3$ -laser, pumped by a $\text{Nd}:\text{YVO}_3$ laser. The laser wavelength was tuned to the D_2 absorption line of cesium, whose resonance wavelength is shifted to 800 nm due to the influence of the surrounding helium matrix. The atomic fluorescence light from the sample volume ($\sim 3 \text{ mm}^3$) is detected by a fiber coupled optical spectrum analyzer (Ando Co. Ltd., AQ-6315A) which has a detection range of 350-1750 nm. The spectral resolution was limited to about 5 nm.

Fig. 1 shows a typical recorded emission spectrum, as well as calculated spectra of several Cs^*He_n exciplexes. The smallest peak at 11400 cm^{-1} corresponds to fluorescence on the D_1 transition, indicating the existence of a transfer channel between $6\text{P}_{3/2}$ and $6\text{P}_{1/2}$ states. That process was studied before, in experiments with superfluid helium [11]. Another emission line appears at 10520 cm^{-1} . We attribute this broad and asymmetric peak to the emission of $\text{Cs}(\text{A}\Pi_{3/2})\text{He}_2$ exciplexes, which had previously been observed in liquid helium and in cold helium gas [3]. The peak presented here is blue-shifted with respect to the one reported in [3] by approximately 500 cm^{-1} . The shift is due to the influence of the helium

*peter.moroshkin@unifr.ch

†www.unifr.ch/physics/frap/




FIG. 1: Measured emission spectrum (dots) of matrix-isolated Cs atoms in the hcp phase of solid ^4He . The fluorescence emission was observed following resonant D_2 excitation of the atoms. For comparison also calculated emission lines (solid curves) of $\text{Cs}(\text{A}\Pi_{3/2})\text{He}_2$, $\text{Cs}(\text{A}\Pi_{1/2})\text{He}_6$, and $\text{Cs}(\text{A}\Pi_{1/2})\text{He}_7$ exciplexes are presented.

matrix and increases with helium pressure. The detailed study of that dependence in the bcc and hcp phases of solid helium will be presented elsewhere [12]. There is yet another prominent emission line at 7130 cm^{-1} , with a linewidth of 1010 cm^{-1} (FWHM), which is almost three times stronger than the former peak. Our theoretical model suggests that this newly discovered intense emission line can be assigned to a higher order ($n > 2$) exciplex.

We have calculated the emission spectra of Cs^*He_n exciplexes with a treatment similar to those of [3, 4]. We consider only the influence of the n helium atoms that form the exciplex and neglect the influence of the bulk of the surrounding helium matrix. The interaction between one cesium atom and the n He atoms is described as the sum over adiabatic molecular two-body interaction potentials. We use the pair potentials between a helium ground state atom and alkali atoms in their ground and lower excited states calculated by Pascale [13]. For the 6S ground state that potential, $V_\sigma^{6s}(r)$, is radially symmetric. For the 6P states the interaction is anisotropic and can be expressed by the operator

$$V^{6P}(\mathbf{r}) = V_\sigma^{6P}(r) + \left(\frac{\mathbf{L} \cdot \mathbf{r}}{\hbar r} \right)^2 [V_\pi^{6P}(r) - V_\sigma^{6P}(r)], \quad (1)$$

where $\mathbf{r} = \mathbf{r}(r, \theta, \varphi)$ denotes the position vector of a helium atom with respect to the cesium atom and $\mathbf{L}$ is the orbital angular momentum operator of the cesium valence electron [1]. Stable exciplexes of the form $\text{Cs}^*\text{He}_{n=2}$ are formed by two helium atoms located on a common axis on opposite sides of the cesium atom. For $\text{Cs}^*\text{He}_{n \geq 3}$ the helium atoms are distributed on a concentric ring around the alkali atom. The summation over the pair

potentials can be expressed by the operator $V_n^{Cs-He}(r) = \sum_{i=1}^n V^{6P}(\mathbf{r}_i)$, with $\mathbf{r}_i = \mathbf{r}(r, \theta = \pi/2, \varphi_i = i2\pi/n)$ and $n = 1, 2, \dots$. In addition we include He-He interactions by summing over the corresponding $V_{He-He}(R)$ potentials between neighboring helium atoms using the semi-empirical potential given by Beck [14]. The distance R between two neighboring helium atoms is a function of the cesium-helium separation r and the number n of helium atoms: $R = |\mathbf{r}_i - \mathbf{r}_{i+1}| = 2r \sin(\pi/n)$. After including the spin-orbit interaction in Cs the total interaction potential of the Cs^*He_n system reads

$$V_{Cs^*\text{He}_n}(r) = V_n^{Cs-He}(r) + nV_{He-He}(R) + 2/3\Delta \mathbf{L} \cdot \mathbf{S}, \quad (2)$$

where $\Delta = 554.0\text{ cm}^{-1}$ is the fine-structure splitting of the free cesium 6P state and $\mathbf{S}$ the electronic spin operator. $V_{Cs^*\text{He}_n}(r)$ is diagonalized algebraically. In Figs. 2 and 3 the resulting r -dependencies of the eigenvalues are shown for Cs^*He_2 and Cs^*He_6 respectively. The same plots also show the ground state potentials given by $nV_\sigma^{6s}(r) + nV_{He-He}(R)$. The potentials are labelled according to their electronic configurations as $X^2\Sigma_{1/2}$, $A^2\Pi_{1/2}$, $A^2\Pi_{3/2}$ and $B^2\Sigma_{1/2}$. The quantization axis is defined by the symmetry axis of the exciplexes, which is the internuclear axis of the cesium atom and the two helium atoms in the case of $\text{Cs}^*\text{He}_{n=1,2}$, whereas for $\text{Cs}^*\text{He}_{n \geq 3}$ it is the axis of the helium ring. Pictographs next to the curves show the variation of the cesium electronic density as the n helium atoms, indicated by two filled circles, approach the cesium atom.

From the adiabatic potentials of Figs. 2 and 3 one sees that the helium atoms, due to the Pauli principle, are repelled by the cesium valence electron. However, in cases where the atoms approach along a nodal line or in a nodal plane of the electron distribution they experience an attractive van der Waals force until they are repelled by the cesium core. This definitely holds for the $\text{Cs}(\text{A}\Pi_{3/2})\text{He}_2$ exciplex. For $\text{Cs}(\text{A}\Pi_{1/2})\text{He}_2$ the situation is more complicated. When the two helium atoms are far away the electronic configuration is the one of the $6\text{P}_{1/2}$ state of the free cesium atom, which has a spherical symmetry and is hence, as the $6\text{S}_{1/2}$ ground state, repulsive for the helium atoms. However, when the helium atoms approach the cesium atom the electronic wave function of the latter is deformed and becomes apple-shaped. This new configuration now offers a binding potential minimum. The formation of this second configuration has a potential barrier of 79 cm^{-1} , which is much higher than the thermal energy of 1.0 cm^{-1} . If, on the other hand, the two atoms do not approach simultaneously, but one after the other, only the first atom has to overcome the potential barrier thereby forming the apple-like electronic configuration of a Cs^*He_1 structure, which then becomes attractive for a second helium atom approaching from the opposite side. A third helium atom approaching the $\text{Cs}(\text{A}\Pi_{1/2})\text{He}_2$ exciplex will again be repelled. However, if it comes sufficiently close, the electronic configuration changes to a dumbbell shape, which has a binding minimum for the three helium atoms. Here too, a potential




FIG. 2: Adiabatic potentials of the Cs^*He_2 system, including the spin-orbit interaction. The two helium atoms are located at r and $-r$ on the quantization (rotationally symmetry) axis, indicated in the pictographs by a solid line. The shape of the electronic density distribution of the cesium atom changes significantly as helium atoms (filled circles) approach.

barrier has to be overcome. The three atoms are then bound and located on a ring concentric around the waist of the dumbbell. The $\text{Cs}(\text{A}\Pi_{1/2})\text{He}_3$ structure exerts a purely attractive potential on further helium atoms. With an increasing number of helium atoms on the ring the repulsive potential between those atoms increases, which puts a natural limit on the maximum number n_{max} that can be accommodated. Only the $\text{A}\Pi_{1/2}$ state can bind more than two helium atoms as can be seen from Fig. 3.

It was pointed out by Dupont-Roc [1] that the height of the potential barrier is determined by the strength of the spin-orbit interaction of the P state. If the L-S coupling is weak compared to the alkali-helium interaction as in the case of Na, K, and Rb, it can be neglected and the electronic configuration can be approximated by a P_z orbital that allows the formation of dumbbell-shaped exciplexes with $n > 2$. If on the other hand the spin-orbit interaction dominates, as for cesium, one has to consider the electron distributions of the L-S-coupled $P_{1/2}$ and $P_{3/2}$ states, which are spherical in the first case and of apple shape in the second case, thereby restricting n_{max} to 2. In contrast to this simple model our treatment takes into account that the electronic configuration changes adiabatically its shape as helium atoms approach.




FIG. 3: Adiabatic potentials of the Cs^*He_6 system. The six helium atoms are located on a ring of radius r concentric with the symmetry axis. Only the $\text{A}^2\Pi_{1/2}$ potential has a binding attractive well.

We calculated the adiabatic potentials and the vibrational zero-point energies for all Cs^*He_n systems up to $n = 9$. The treatment of the molecular vibrations is approximate and will be described elsewhere. Higher vibrational states are not populated at the temperature of the experiments and can be neglected, as well as the contributions from rotations.

The n dependence of the zero-point energies for $\text{Cs}(\text{A}\Pi_{1/2})\text{He}_n$ exciplexes is represented in Fig. 4 by filled circles. Also shown are the heights of the potential barriers (open squares) and the depths of the potential minima (open circles). All energies are given with respect to the dissociation limit, i.e., the electronic energy of the $6P_{1/2}$ state. The significant increase of the potential barrier when going from $n = 2$ to $n = 3$ reflects the discussion given above. The configuration with $n = 8$ has the deepest attractive potential, while its zero-point energy due to the stronger localization of helium atoms exceeds the barrier energy, which makes that complex unstable. $\text{Cs}(\text{A}\Pi_{1/2})\text{He}_7$ seems to be the largest possible quasi-bound exciplex. Only the molecule with $n = 6$ has an energy below the dissociation limit and is hence predicted to be stable. Note that for rubidium several configurations are predicted to be stable and $\text{Rb}(\text{A}\Pi_{1/2})\text{He}_n$ exciplexes with $n = 1 \dots 6$ have been observed recently [4].

In a last step we calculated the shape of the emis-




FIG. 4: Energy dependencies of the $\text{Cs}(A^2\Pi_{1/2})\text{He}_n$ exciplexes as a function of the number n of helium atoms. Shown are the minimal energies (open circles) of the potential wells, the barrier heights (open squares) and the total zero-point energies (filled circles). All energies are given with respect to the dissociation limit, i.e., the electronic energy of the $6P_{1/2}$ state. Corresponding points are joined by lines to guide the eye. The temperature of the helium matrix corresponds to $kT = 1.0 \text{ cm}^{-1}$.

sion lines by using the Franck-Condon approximation in a similar way as the authors of [3]. Among all exciplexes considered, the best agreement of the calculated spectra with the experimental ones is provided by those shown in Fig. 1 by solid curves. The shape and width of the line at 10520 cm^{-1} agrees well with the calculated emission line of $\text{Cs}(A\Pi_{3/2})\text{He}_2$. However, the apparently excellent agreement between the positions of these lines is accidental as the pressure shift, i.e., the influence of the surrounding helium matrix was not taken into account. In fact, K. Enomoto et al. [3] measured the emission line of $\text{Cs}(A\Pi_{3/2})\text{He}_2$ exciplexes in cold helium vapor environment to lie at about 10000 cm^{-1} . We like to mention that we have also observed fluorescence from $\text{Cs}(A\Pi_{1/2})\text{He}_2$ exciplexes, not discussed here, following D_1 excitation [12, 15]. Such exciplexes were also observed

recently on cesium doped He clusters [16].

The correct assignment of the strongest measured peak at 7130 cm^{-1} in Fig. 1 is a more subtle task. It is clear that the line originates from a $\text{Cs}(A\Pi_{1/2})\text{He}_{n>2}$ exciplex, in which the helium atoms form a ring in the nodal plane of the Cs atom. It is very likely that this line also shifts with helium pressure. Moreover, as the ring-shaped exciplexes contain a larger number of helium atoms any imprecision of the initial pair potentials will be amplified. As a consequence the line positions of the higher order Cs^*He_n systems cannot be predicted with sufficient accuracy to allow an unambiguous assignment of the observed line. The overall good agreement between calculated and measured lineshapes indicates that the line at 7130 cm^{-1} originates from the decay of an exciplex with a specific number n of bound helium atoms and that it is not a superposition of lines from exciplexes with different values of n . Such superpositions were observed in the case of Rb^*He_n exciplexes in cold ^4He vapor [4]. The theoretical considerations suggest, as a best guess, that the observed peak at 7130 cm^{-1} originates from decaying Cs^*He_6 complexes, as those exciplexes have minimal energy and are the only ones predicted to be stable.

In summary, we have performed a study of laser induced fluorescence of cesium atoms trapped in the hcp phase of a helium crystal. We have reported the observation of new spectral features, which are broader and more intense than the pure atomic lines. We believe that those lines are formed by the emission from two types of specific Cs^*He_n exciplex structures, viz., an apple-shaped complex with two helium atoms bound to the Cs atom and a dumbbell-shaped complex, in which a ring of helium atoms is bound to the nodal plane of the Cs wave function. These assignments are supported by model calculations, which allow us to obtain the corresponding emission spectra. In the case of the ring structure the calculations suggest $n = 6$ as the most likely number of bound atoms.

We like to thank J. Pascale for sending us his numerical Cs-He pair potentials. This work was supported by a grant of the Schweizerischer Nationalfonds.

-
- [1] J. Dupont-Roc, Z. Phys. B **98**, 383 (1995).
 - [2] S. Kanorsky, A. Weis, M. Arndt, R. Dziewior, and T. W. Hänsch, Z. Phys. B **98**, 371 (1995).
 - [3] K. Enomoto, K. Hirano, M. Kumakura, Y. Takahashi, and T. Yabuzaki, Phys. Rev. A **66**, 042505 (2002).
 - [4] K. Hirano, K. Enomoto, M. Kumakura, Y. Takahashi, and T. Yabuzaki, Phys. Rev. A **68**, 012722 (2003).
 - [5] F. Stienkemeier, J. Higgins, C. Callegari, S. I. Kanorsky, W. E. Ernst, and G. Scoles, Z. Phys. D **38**, 253 (1996).
 - [6] J. Reho, J. Higgins, C. Callegari, K. K. Lehmann, and G. Scoles, J. Chem. Phys. **113**, 9686 (2000).
 - [7] F. R. Bruhl, R. A. Trasca, and W. E. Ernst, J. Chem. Phys. **115**, 10220 (2001).
 - [8] C. Schulz, P. Claas, and F. Stienkemeier, Phys. Rev. Lett. **87**, 153401 (2001).
 - [9] S. I. Kanorsky, M. Arndt, R. Dziewior, A. Weis, and T. W. Hänsch, Phys. Rev. B **49**, 3645 (1994).
 - [10] D. Nettels, R. Müller-Siebert, S. Ulzega, and A. Weis, Applied Physics B **77**, 563 (2003).
 - [11] T. Kinoshita, K. Fukuda, Y. Takahashi, and T. Yabuzaki, Phys. Rev. A **52**, 2707 (1995).
 - [12] D. Nettels, A. Hofer, P. Moroshkin, R. Müller-Siebert, S. Ulzega, and A. Weis, to be submitted.
 - [13] J. Pascale, Phys. Rev. A **28**, 632 (1983).
 - [14] D. E. Beck, Molecular Physics **14**, 311 (1968).
 - [15] D. Nettels, Ph.D. thesis, University Fribourg, Switzerland (2003).
 - [16] F. Stienkemeier, private communication.